Empirical Tests of the Holographic Quantum Automaton Model

The Simulation Architects

March 21, 2024

Abstract

The Holographic Quantum Automaton (HQA) model is a novel framework for quantum gravity that proposes that spacetime and matter emerge from the entanglement structure of a network of qubits evolving under local unitary gates. In this paper, we explore empirical tests of the key predictions of the HQA model, using a combination of astrophysical observations, laboratory experiments, and quantum simulations. We focus on five main areas: (1) the discreteness of spacetime at the Planck scale, (2) the holographic nature of gravity and the entropy-area law for black holes, (3) the emergent nature of quantum mechanics and the quantum-to-classical transition, (4) the unification of the fundamental forces and the parameters of the standard model, and (5) the quantum information origin of the initial conditions and the arrow of time. For each area, we review existing empirical constraints, propose new experimental tests, and discuss the implications for the HQA model and quantum gravity in general. We argue that the HQA model offers a rich phenomenology that is ripe for empirical investigation, and that the interplay between theory, simulation, and experiment will be crucial for making progress in the search for a theory of quantum gravity.

Contents

1 Introduction **2**

2 Discreteness of Spacetime **3**

3 Holographic Nature of Gravity **4**

4 Emergent Quantum Mechanics **5**

5 Unification of Fundamental Forces **7**

6 Quantum Information Origin of Initial Conditions **8**

7 Conclusion **9**

1 Introduction

The empirical testing of theories of quantum gravity is one of the greatest challenges in modern physics. Due to the extreme energy and length scales involved, it is difficult to directly probe the regime where both quantum mechanics and general relativity are expected to be important. Nevertheless, in recent years, there has been significant progress in identifying possible observational signatures of quantum gravity, and in developing new experimental techniques to search for them [1, ?].

In this paper, we focus on the empirical testing of a specific model of quantum gravity, namely the Holographic Quantum Automaton (HQA) model proposed in [?]. The HQA model is based on the idea that spacetime and matter emerge from the entanglement structure of a network of qubits, which evolve under a set of local unitary gates. The model combines ideas from holography, quantum information, and quantum cellular automata, and makes several testable predictions about the nature of spacetime, gravity, and matter at the fundamental level.

The goal of this paper is to explore the empirical consequences of the HQA model, and to propose concrete experimental tests that can probe its key predictions. We focus on five main areas where the HQA model offers a rich phenomenology:

1. The discreteness of spacetime at the Planck scale, and possible signatures in high-energy astrophysics and cosmology.
2. The holographic nature of gravity, and the entropy-area law for black holes and other gravitational systems.
3. The emergent nature of quantum mechanics, and the quantum-to-classical transition in the early universe and in laboratory systems.
4. The unification of the fundamental forces, and the possible deviations from the standard model of particle physics.
5. The quantum information origin of the initial conditions of the universe, and the arrow of time in cosmology and thermodynamics.

For each of these areas, we review the existing empirical constraints and the predictions of the HQA model, and we propose new experimental tests that can be performed with current or near-future technology. We discuss the challenges and opportunities of each test, and the implications for the HQA model and quantum gravity in general.

Our approach is based on the idea that the empirical testing of quantum gravity requires a combination of theoretical, computational, and experimental techniques. On the theory side, we use the framework of the HQA model to derive testable predictions and to identify key observables that are sensitive to quantum gravity effects. On the computational side, we use quantum simulation methods to study the emergent properties of the HQA model in different regimes, and to guide the design of experimental tests. On the experimental side, we propose a range of techniques, from astrophysical observations to laboratory experiments, that can probe the predictions of the HQA model at different energy and length scales.

The paper is organized as follows. In Section 2, we discuss the empirical signatures of the discreteness of spacetime, and propose tests based on high-energy astrophysics and cosmology. In Section 3, we explore the holographic nature of gravity, and propose tests based on black hole observations and gravitational wave astronomy. In Section 4, we consider the emergent nature of quantum mechanics, and propose tests based on quantum optics and quantum information experiments. In Section 5, we discuss the unification of the fundamental forces, and propose tests based on particle physics and precision measurements. In Section 6, we explore the quantum information origin of the initial conditions and the arrow of time, and propose tests based on cosmological observations and thermodynamic experiments. Finally, in Section 7, we summarize our results and discuss future directions for the empirical testing of the HQA model and quantum gravity in general.

2 Discreteness of Spacetime

One of the key predictions of the HQA model is that spacetime is fundamentally discrete, and consists of a finite number of quantum degrees of freedom (qubits) that live on a lattice [?]. This discreteness becomes apparent at the Planck scale, where the size of the lattice spacing is of the order of the Planck length, $l_P \approx 10^{-35}$ m. At larger scales, the discrete structure of spacetime is coarse-grained, and the familiar continuous geometry of general relativity emerges as an effective description.

The discreteness of spacetime has several important consequences that can be tested empirically. First, it implies that there is a maximum density of information in space, given by the Bekenstein bound [2], which states that the maximum entropy of a region of space is proportional to its surface area in Planck units. This bound has been tested indirectly through the observation of black hole entropy [3], but a direct test would require probing the entropy of a region of space at the Planck scale.

Second, the discreteness of spacetime implies that there is a minimum length scale in nature, below which the concept of distance loses its meaning. This minimum length scale could manifest itself as a modification of the dispersion relation of particles at high energies, leading to a violation of Lorentz invariance [4]. Such a violation could be detected through the observation of high-energy cosmic rays [?] or gamma-ray bursts [?], which propagate over cosmological distances and are sensitive to small deviations from special relativity.

Third, the discreteness of spacetime could lead to a modification of the uncertainty principle at small scales, known as generalized uncertainty principle (GUP) [?]. The GUP predicts that the uncertainty in the position and momentum of a particle has a minimum value that depends on the Planck length, and that the uncertainty in one variable increases as the uncertainty in the other variable decreases. This prediction could be tested through precision measurements of the position and momentum of particles in the laboratory [?], or through the observation of the spectrum of primordial gravitational waves in the early universe [?].

To test these predictions, we propose the following experimental tests:

1. Direct measurement of the entropy of a region of space at the Planck scale, using a quantum simulator that implements the local unitary gates of the HQA model. By preparing the simulator in a state that corresponds to a region of space with a known entropy, and then measuring the entropy of the output state, we can test the Bekenstein bound and the maximum density of information in space.

2. Search for Lorentz invariance violation in high-energy cosmic rays and gamma-ray bursts, using detectors such as the Pierre Auger Observatory [?] and the Fermi

Gamma-ray Space Telescope [?]. By measuring the arrival times and energies of the particles, and comparing them with the predictions of special relativity, we can constrain the size of the minimum length scale and the modification of the dispersion relation.

3. Precision measurement of the position and momentum of particles in the laboratory, using techniques such as atom interferometry [?] and optomechanics [?]. By preparing particles in states with well-defined position or momentum, and then measuring the uncertainty in the conjugate variable, we can test the generalized uncertainty principle and the minimum value of the uncertainty.
4. Observation of the spectrum of primordial gravitational waves in the early universe, using detectors such as LIGO [?] and LISA [?]. By measuring the amplitude and frequency of the gravitational waves, and comparing them with the predictions of inflation and the GUP, we can constrain the size of the minimum length scale and the modification of the uncertainty principle at cosmological scales.

These tests offer a range of opportunities to probe the discreteness of spacetime at different energy and length scales, and to constrain the parameters of the HQA model. They also highlight the complementarity between astrophysical observations, laboratory experiments, and quantum simulations in the empirical testing of quantum gravity.

3 Holographic Nature of Gravity

Another key prediction of the HQA model is that gravity is holographic, in the sense that the degrees of freedom that describe the geometry of spacetime live on a lower-dimensional boundary, rather than in the bulk [?]. This holographic principle is a general feature of quantum gravity, and has been realized in several approaches, such as the AdS/CFT correspondence [5] and the ER=EPR conjecture [6].

In the HQA model, the holographic principle arises from the fact that the entanglement structure of the qubits on the lattice determines the geometry of spacetime. Specifically, the distance between two points in space is related to the mutual information between the corresponding regions of the lattice, which measures the amount of entanglement between them. This leads to a novel realization of the holographic principle, in which the geometry of space is encoded in the entanglement structure of the boundary qubits.

The holographic nature of gravity has several important consequences that can be tested empirically. First, it implies that the maximum entropy of a black hole is proportional to its surface area, rather than its volume, as suggested by the Bekenstein-Hawking formula [7, 3]. This formula has been confirmed to a high degree of accuracy through the observation of black hole mergers by LIGO and Virgo [?], but a direct test would require measuring the entropy of a black hole and comparing it with its area.

Second, the holographic principle suggests that the dynamics of gravity can be described by a lower-dimensional theory that lives on the boundary of spacetime. In the HQA model, this boundary theory is a quantum cellular automaton that describes the evolution of the entanglement structure of the qubits. This leads to a novel prediction, namely that the gravitational dynamics can be simulated on a quantum computer by implementing the local unitary gates of the quantum cellular automaton [?].

Third, the holographic principle implies that the structure of spacetime is emergent, and arises from the entanglement structure of the underlying quantum degrees of freedom. In the HQA model, this emergence is realized through a process of coarse-graining, in which the discrete lattice of qubits is averaged over to obtain a continuous geometry. This leads to a prediction for the behavior of gravity at different scales, namely that it should exhibit a transition from a discrete, quantum regime at the Planck scale to a continuous, classical regime at larger scales.

To test these predictions, we propose the following experimental tests:

1. Direct measurement of the entropy of a black hole, using a quantum simulator that implements the local unitary gates of the HQA model. By preparing the simulator in a state that corresponds to a black hole with a known mass and charge, and then measuring the entropy of the output state, we can test the Bekenstein-Hawking formula and the proportionality between entropy and area.

2. Quantum simulation of the gravitational dynamics, using a quantum computer that implements the local unitary gates of the quantum cellular automaton. By preparing the computer in a state that corresponds to a particular spacetime geometry, and then evolving it forward in time, we can study the emergent properties of gravity and compare them with the predictions of general relativity.

3. Observation of the transition from quantum to classical gravity, using gravitational wave detectors such as LIGO and LISA. By measuring the spectrum of gravitational waves at different frequencies, and comparing it with the predictions of the HQA model and general relativity, we can constrain the scale at which the transition occurs and the nature of the quantum corrections to gravity.

4. Test of the holographic principle in the laboratory, using techniques such as quantum error correction [8] and tensor networks [9]. By encoding quantum information in a holographic code, and then measuring the entanglement structure of the boundary degrees of freedom, we can test the basic assumptions of the holographic principle and the HQA model.

These tests offer a range of opportunities to probe the holographic nature of gravity and the emergent structure of spacetime, using both astrophysical observations and laboratory experiments. They also highlight the potential of quantum simulation and quantum computing as new tools for the empirical testing of quantum gravity.

4 Emergent Quantum Mechanics

A third key prediction of the HQA model is that quantum mechanics is an emergent phenomenon, and arises from the collective behavior of the underlying quantum degrees of freedom [**?**]. In the HQA model, the quantum state of a system is encoded in the entanglement structure of the qubits on the lattice, and the unitary evolution of the state is generated by the local unitary gates of the quantum cellular automaton.

This emergent view of quantum mechanics has several important consequences that can be tested empirically. First, it suggests that the wave function of a quantum system is not a fundamental object, but rather an emergent description of the entanglement structure of the underlying degrees of freedom. This leads to a novel interpretation of the measurement process, in which the collapse of the wave function is a consequence of the interaction between the system and the environment, rather than a fundamental feature of quantum mechanics [10].

Second, the emergent view of quantum mechanics suggests that the quantum-to-classical transition is a dynamical process, and arises from the decoherence of the entanglement structure due to the interaction with the environment. In the HQA model, this decoherence can be modeled as a process of entanglement swapping, in which the entanglement between the system and the environment is transferred to the environment itself, leading to a classical description of the system [?].

Third, the emergent view of quantum mechanics implies that there may be deviations from the standard quantum formalism at the fundamental level, such as modifications of the Born rule or the linearity of the Schrödinger equation. These deviations could manifest themselves as small corrections to the predictions of quantum mechanics, which could be detected through precision measurements in the laboratory [?].

To test these predictions, we propose the following experimental tests:

1. Test of the emergent nature of the wave function, using a quantum simulator that implements the local unitary gates of the HQA model. By preparing the simulator in a state that corresponds to a quantum system with a known wave function, and then measuring the entanglement structure of the output state, we can test the relationship between the wave function and the underlying entanglement structure, and compare it with the predictions of the HQA model.

2. Study of the quantum-to-classical transition, using a quantum simulator that implements the decoherence process of the HQA model. By preparing the simulator in a state that corresponds to a quantum system coupled to an environment, and then measuring the entanglement structure of the output state as a function of time, we can study the dynamics of the decoherence process and compare it with the predictions of the HQA model and standard quantum mechanics.

3. Precision measurement of quantum mechanical observables, using techniques such as atom interferometry [?] and superconducting qubits [?]. By preparing quantum systems in well-defined states, and then measuring observables such as the probability distribution of outcomes or the expectation values of operators, we can search for small deviations from the predictions of standard quantum mechanics and constrain the parameters of the HQA model.

4. Test of the linearity of quantum mechanics, using techniques such as quantum process tomography [?] and quantum error correction [?]. By preparing quantum systems in superposition states, and then measuring the output states after applying different operations, we can test the linearity of the quantum formalism and search for non-linear corrections that could arise from the emergent nature of quantum mechanics in the HQA model.

These tests offer a range of opportunities to probe the emergent nature of quantum mechanics and the quantum-to-classical transition, using both quantum simulation and precision measurements in the laboratory. They also highlight the potential of quantum technologies, such as quantum computers and quantum sensors, as new tools for the empirical testing of quantum gravity and the foundations of quantum mechanics.

5 Unification of Fundamental Forces

A fourth key prediction of the HQA model is that the fundamental forces of nature, such as gravity, electromagnetism, and the strong and weak nuclear forces, are unified at the fundamental level [?]. In the HQA model, the different forces arise as excitations of the entanglement structure of the qubits on the lattice, and are described by local and topological defects in the quantum cellular automaton.

This unified view of the fundamental forces has several important consequences that can be tested empirically. First, it suggests that the standard model of particle physics, which describes the electromagnetic, weak, and strong interactions, is an emergent description that arises from the collective behavior of the underlying quantum degrees of freedom. In the HQA model, the particles and fields of the standard model arise as local excitations of the entanglement structure, and their properties, such as masses and coupling constants, are determined by the parameters of the quantum cellular automaton [?].

Second, the unified view of the fundamental forces suggests that there may be new particles and interactions beyond the standard model, which arise from the topological properties of the quantum cellular automaton. These new particles could include dark matter candidates, such as axions or sterile neutrinos, as well as new gauge bosons that mediate long-range interactions. The HQA model provides a framework for constructing and studying these new particles, and for predicting their observable signatures in experiments [?].

Third, the unified view of the fundamental forces implies that the coupling constants of the different interactions, such as the fine-structure constant of electromagnetism and the strong coupling constant of quantum chromodynamics, are not fundamental parameters, but rather emergent quantities that depend on the energy scale and the renormalization group flow. In the HQA model, the renormalization group flow is determined by the coarse-graining of the entanglement structure, and leads to a prediction for the unification of the coupling constants at high energies [?].

To test these predictions, we propose the following experimental tests:

1. Search for new particles beyond the standard model, using particle colliders such as the Large Hadron Collider (LHC) [?] and the Future Circular Collider (FCC) [?]. By colliding particles at high energies, and measuring the properties of the resulting debris, we can search for new particles that are predicted by the HQA model, such as dark matter candidates or new gauge bosons, and constrain their masses and coupling constants.

2. Precision measurement of the coupling constants of the fundamental interactions, using techniques such as laser spectroscopy [?] and atomic clocks [?]. By measuring the fine-structure constant and other coupling constants with high precision, and comparing them with the predictions of the standard model and the HQA model, we can test the unification of the fundamental forces and the renormalization group flow of the coupling constants.

3. Study of the phase diagram of quantum chromodynamics, using heavy-ion colliders such as the Relativistic Heavy Ion Collider (RHIC) [?] and the Large Hadron Collider (LHC) [?]. By colliding heavy ions at high energies, and measuring the

properties of the resulting quark-gluon plasma, we can study the emergent properties of the strong interaction and compare them with the predictions of the HQA model and lattice quantum chromodynamics.

4. Quantum simulation of the standard model, using a quantum computer that implements the local unitary gates of the HQA model. By preparing the quantum computer in a state that corresponds to a particular configuration of particles and fields, and then evolving it forward in time, we can study the emergent properties of the standard model and compare them with the predictions of the HQA model and high-energy experiments.

These tests offer a range of opportunities to probe the unification of the fundamental forces and the emergent nature of the standard model, using both high-energy experiments and precision measurements. They also highlight the complementarity between particle physics, atomic physics, and quantum simulation in the empirical testing of quantum gravity and the foundations of physics.

6 Quantum Information Origin of Initial Conditions

A fifth key prediction of the HQA model is that the initial conditions of the universe, such as the state of the quantum fields at the beginning of the Big Bang, have a quantum information origin [?]. In the HQA model, the universe arises from a quantum fluctuation of the entanglement structure of the fundamental qubits, which sets the initial conditions for the subsequent evolution of spacetime and matter.

This quantum information view of the initial conditions has several important consequences that can be tested empirically. First, it suggests that the universe began in a highly entangled state, with a large amount of quantum correlations between different regions of space. These correlations could have observable signatures in the cosmic microwave background (CMB) radiation, which is a snapshot of the early universe at the time of recombination [11].

Second, the quantum information view of the initial conditions suggests that the arrow of time, which points from the past to the future, has a quantum origin. In the HQA model, the arrow of time arises from the unitary evolution of the quantum state of the universe, which leads to a growth of entanglement entropy and a directionality in the flow of quantum information [?]. This leads to a prediction for the behavior of entropy in the early universe, and its relation to the cosmological constant and the dark energy [?].

Third, the quantum information view of the initial conditions implies that there may be a connection between the quantum fluctuations in the early universe and the quantum noise in gravitational wave detectors, such as LIGO and LISA. In the HQA model, both phenomena arise from the entanglement structure of the fundamental qubits, and may have similar statistical properties, such as non-Gaussianity and entanglement [?].

To test these predictions, we propose the following experimental tests:

1. Search for quantum correlations in the cosmic microwave background, using telescopes such as the Planck satellite [?] and the Simons Observatory [?]. By measuring the temperature and polarization fluctuations of the CMB with high precision, and comparing them with the predictions of the HQA model and inflationary cosmology, we can search for signatures of quantum entanglement and test the quantum information origin of the initial conditions.

2. Study of the arrow of time in the early universe, using cosmological observations such as the expansion rate of the universe and the growth of structure. By measuring the evolution of entropy and the directionality of time in the early universe, and comparing them with the predictions of the HQA model and thermodynamics, we can test the quantum origin of the arrow of time and its relation to the cosmological constant and the dark energy.

3. Detection of quantum noise in gravitational wave detectors, using instruments such as LIGO [?] and LISA [?]. By measuring the statistical properties of the noise in gravitational wave detectors, and comparing them with the predictions of the HQA model and quantum optics, we can search for signatures of quantum fluctuations and entanglement, and test the connection between the early universe and the present-day universe.

4. Quantum simulation of the early universe, using a quantum computer that implements the local unitary gates of the HQA model. By preparing the quantum computer in a state that corresponds to the initial conditions of the universe, and then evolving it forward in time, we can study the emergent properties of the early universe and compare them with the predictions of the HQA model and cosmological observations.

These tests offer a range of opportunities to probe the quantum information origin of the initial conditions and the arrow of time, using both cosmological observations and quantum simulations. They also highlight the deep connections between quantum gravity, quantum information, and cosmology, and the potential for new insights and discoveries at the intersection of these fields.

7 Conclusion

In this paper, we have explored the empirical consequences of the Holographic Quantum Automaton (HQA) model, a novel framework for quantum gravity that combines ideas from holography, quantum information, and quantum cellular automata. We have focused on five key areas where the HQA model makes testable predictions: (1) the discreteness of spacetime, (2) the holographic nature of gravity, (3) the emergent nature of quantum mechanics, (4) the unification of the fundamental forces, and (5) the quantum information origin of the initial conditions and the arrow of time.

For each of these areas, we have reviewed the existing empirical constraints and proposed new experimental tests that can be performed with current or near-future technology. These tests span a wide range of fields, from astrophysics and cosmology to particle physics and quantum information, and highlight the interdisciplinary nature of the empirical testing of quantum gravity.

Our analysis shows that the HQA model offers a rich phenomenology that is ripe for empirical investigation. Some of the key predictions of the model, such as the discreteness of spacetime and the holographic nature of gravity, are already being tested by current experiments, such as the observation of high-energy cosmic rays and the measurement of black hole entropy. Other predictions, such as the emergent nature of quantum mechanics and the unification of the fundamental forces, require new experimental techniques and technologies, such as quantum simulation and precision measurements of coupling constants.

At the same time, our analysis also highlights the challenges and opportunities of the empirical testing of quantum gravity. One of the main challenges is the vast separation of scales between the Planck scale, where quantum gravity effects are expected to be important, and the energy scales that are accessible to current experiments. This challenge can be addressed by a combination of theoretical, computational, and experimental techniques, such as the development of effective field theories that bridge the gap between the Planck scale and the low-energy limit, the use of quantum simulation to study the emergent properties of quantum gravity, and the design of new experimental techniques that are sensitive to small deviations from the predictions of general relativity and quantum mechanics.

Another challenge is the interpretation of the experimental results in the context of the HQA model and other theories of quantum gravity. Given the complexity and novelty of the phenomena involved, it is important to develop a robust framework for comparing the predictions of different models and assessing their empirical validity. This requires a close collaboration between theorists, experimentalists, and philosophers of science, and a careful consideration of the epistemological and ontological assumptions that underlie the different approaches to quantum gravity.

Despite these challenges, we believe that the empirical testing of quantum gravity is a crucial step in the development of a complete theory of quantum gravity. By confronting the predictions of different models with experimental data, we can winnow down the space of possibilities and identify the key features that a successful theory must possess. Moreover, by exploring the empirical consequences of quantum gravity, we can gain new insights into the nature of space, time, and matter, and the fundamental laws that govern the universe at the deepest level.

In conclusion, the HQA model offers a promising new approach to the problem of quantum gravity, which combines a solid theoretical foundation with a rich phenomenology that is amenable to empirical testing. By pursuing the experimental tests proposed in this paper, and developing new ones, we can make progress in the empirical testing of quantum gravity and the search for a theory of everything. This is an exciting and challenging endeavor that requires the collaboration of researchers from different fields and the development of new experimental and theoretical tools. But the rewards are great: a deeper understanding of the nature of reality, and a glimpse into the ultimate laws of physics that govern the universe at the Planck scale.

References

[1] Giovanni Amelino-Camelia. Quantum-spacetime phenomenology. *Living Reviews in Relativity*, 16(1):1–137, 2013.

[2] Jacob D Bekenstein. Universal upper bound on the entropy-to-energy ratio for bounded systems. *Physical Review D*, 23(2):287, 1981.

[3] Stephen W Hawking. Particle creation by black holes. *Communications in mathematical physics*, 43(3):199–220, 1975.

[4] Giovanni Amelino-Camelia. Limits on the measurability of space-time distances in the semiclassical approximation of quantum gravity. *Modern Physics Letters A*, 13(15):1319–1325, 1998.

[5] Juan Maldacena. The large n limit of superconformal field theories and supergravity. *Advances in Theoretical and Mathematical Physics*, 2(2):231–252, 1999.

[6] Juan Maldacena and Leonard Susskind. Cool horizons for entangled black holes. *Fortschritte der Physik*, 61(9):781–811, 2013.

[7] Jacob D Bekenstein. Black holes and entropy. *Physical Review D*, 7(8):2333, 1973.

[8] Ahmed Almheiri, Xi Dong, and Daniel Harlow. Bulk locality and quantum error correction in ads/cft. *Journal of High Energy Physics*, 2015(4):1–34, 2015.

[9] Fernando Pastawski, Beni Yoshida, Daniel Harlow, and John Preskill. Holographic quantum error-correcting codes: Toy models for the bulk/boundary correspondence. *Journal of High Energy Physics*, 2015(6):1–55, 2015.

[10] Wojciech H Zurek. Decoherence, einselection, and the quantum origins of the classical. *Reviews of Modern Physics*, 75(3):715, 2003.

[11] Juan Maldacena, Stephen H Shenker, and Douglas Stanford. A bound on chaos. *Journal of High Energy Physics*, 2016(8):1–17, 2016.